


## Emergency Treatment of Opioid Overdose

### Important:

- For use in the nose only
- Do not test nasal spray device before use
- 1 nasal spray device contains 1 dose of medicine
- Each device sprays 1 time only



### Drug Facts

#### Active ingredient (in each spray)

Naloxone hydrochloride 4 mg..... Emergency treatment of opioid overdose

#### Purpose

#### Uses


- to “revive” someone during an overdose from many **prescription pain medications** or **street drugs such as heroin**
- this medicine can save a life

#### Directions



#### 1 CHECK ▶ Step 1: CHECK if you suspect an overdose:

- **CHECK** for a suspected overdose: the person will not wake up or is very sleepy or not breathing well
  - yell “Wake up!”
  - shake the person gently
  - if the person is not awake, go to Step 2



#### 2 GIVE ▶ Step 2: Give 1st dose in the nose

- **HOLD** the nasal spray device with your thumb on the bottom of the plunger
- **INSERT** the nozzle into either NOSTRIL
- **PRESS** the plunger firmly to give the 1st dose
- 1 nasal spray device contains 1 dose



#### 3 CALL ▶ Step 3: Call 911

- **CALL 911** immediately after giving the 1<sup>st</sup> dose



#### 4 WATCH/GIVE ▶ Step 4: WATCH & GIVE

- **WAIT** 2-3 minutes after the 1<sup>st</sup> dose to give the medicine time to work
- if the person wakes up: Go to Step 5
- if the person does not wake up:
  - **CONTINUE TO GIVE** doses every 2-3 minutes until the person wakes up
  - it is safe to keep giving doses



#### 5 STAY ▶ Step 5: STAY

- **STAY** until ambulance arrives: even if the person wakes up
- **GIVE** another dose if the person becomes very sleepy again
- You may need to give all the doses in the pack

Unfold for  
Drug Facts

Unfold for  
Drug Facts



Drug Facts  
Continued on  
Other Side ▶

## **Drug Facts** (continued)

---

### **Warning**

**When using this product** some people may experience symptoms when they wake up, such as shaking, sweating, nausea, or feeling angry. This is to be expected.

### **Other information**

- store at room temperature or refrigerator, between 2°C to 25°C (36°F to 77°F)
- do not freeze
- avoid excessive heat above 40°C (104°F)
- protect from light
- the product is packaged in individually-sealed blisters. Do not use if the blister is open or torn, or if the device appears damaged.

### **Inactive ingredients**

benzalkonium chloride, edetate disodium, hydrochloric acid, sodium chloride, water

### **Questions?**

call 1-844-4NARCAN (1-844-462-7226) or go to [www.narcan.com](http://www.narcan.com)